

COLONOSCOPY- EXTENDED GOLYTELY BOWEL PREPARATION



Date & Location

You should have received the date and location of your colonoscopy from our practice by phone, through the portal, and/or email and text messaging. If you do not know your procedure date or location, please contact our practice. Your procedure time will be communicated to you by the facility the day before your procedure.



Shopping List

- GoLYTELY or (generic name GaviLyle) prescription bowel preparation prescribed to your pharmacy (2 kits)
- Four (4) 5 milligram tablets of Dulcolax (or generic named bisacodyl)
- Four (4) 80 milligram tablets of simethicone or two (2) 250 milligram tablets of Phazyme



Medications

If you take any of the following medications, please inform our scheduling staff as soon as possible so we can coordinate stopping these medications with your prescribing physician:

- **Blood thinners** including but not limited to Coumadin (warfarin), Plavix (clopidogrel), Xarelto (rivaroxaban), Eliquis (apixaban), Brilinta (tricagrelor), Pradaxa (dabigartran).
- **GLP1 medications** such as Trulicity (dulaglutide), Ozempic (semaglutide), Victoza (liraglutide), Bydureon (exenatide) used for diabetes or weight loss.



Transportation

Driving is not permitted until the day after your procedure. Arrange for a ride to and from the center from someone 18 or older who can take responsibility for your care. That person should arrive no later than 1 hour after your procedure time.



Cancellation/No-Show

Our practice charges \$100 for a colonoscopy and \$200 for a combined colonoscopy/EGD that is cancelled within 7 business days of a procedure. For patients who have canceled in the past, a \$100 refundable deposit will be taken to secure your appointment. The deposit will be refunded after your procedure is completed if your account is in good standing.

On Page 2 is a calendar to walk you through your prep which starts 7 days before your procedure date. You can find details regarding low residue foods and clear liquids on our website at

<https://bedfordgastroenterology.com/colonoscopy-prep-faq/>

Please feel free to call us with any questions you may have about these instructions

Calendar for Extended GoLYTELY Bowel Preparation

7 Days or More Before Procedure	5 Days Before Procedure	2 Days Before Procedure	1 Day Before Procedure	Day of Procedure
<ul style="list-style-type: none"> ❑ Make sure you've completed your shopping list. ❑ Contact your insurance company with benefit questions. ❑ If you take iron supplements, stop taking them 7 days before your procedure. ❑ If you take insulin, ask your prescribing physician (primary care or endocrinologist) for instructions on how much insulin to take 1 day before and on the day of the procedure. ❑ If your location is Bedford Ambulatory Surgery Center or Surgical Center of NH, complete your health history in One Medical Passport (sent by facility). ❑ To cancel without incurring a fee, you must call our office at least 7 days before the procedure. 	<ul style="list-style-type: none"> ❑ Eat only low residue foods from this day until after your procedure. Examples of low residue foods can be found on our website as indicated on page 1. ❑ Confirm your ride to and from the procedure as per page 1. 	<ul style="list-style-type: none"> ❑ You may have a <u>light</u> breakfast and lunch before 1PM. Do not eat dinner. Continue to avoid high residue foods. ❑ Add water to the fill line of 1 of your GoLYTELY containers and shake to dissolve the powder into the mixture. Store in refrigerator. ❑ Drink all of the 4 liter mixture by consuming 8 ounces every 15 to 20 minutes. If you develop nausea, drink more slowly and try walking as you drink. 	<ul style="list-style-type: none"> ❑ Drink a clear liquid diet the entire day. Do not consume dairy or solid food on this day including breakfast. ❑ Examples of clear liquids can be found on our website as indicated on page 1. ❑ Take two Dulcolax (bisacodyl) tablets between 3PM and 6PM. Take <u>either</u> two 80 mg simethicone tablets <u>or</u> one 250 mg Phazyme tablet. ❑ Add water to the fill line of your 2nd GoLYTELY container and shake to dissolve the powder into the mixture. Store in refrigerator. ❑ Drink one half (2 liters) of the mixture consuming 8 ounces every 15 to 20 minutes. If you develop nausea, drink more slowly and try walking as you drink. Save the remaining half for the next day. 	<ul style="list-style-type: none"> ❑ Take two 80 mg simethicone tablets <u>or</u> one 250 mg Phazyme tablet 5 to 6 hours before your arrival time. ❑ Take two Dulcolax (bisacodyl) tablets immediately after the simethicone or Phazyme tablets. ❑ Drink the remaining half of the GoLYTELY mixture consuming 8 ounces every 15 to 20 minutes until completely gone. ❑ Do not drink any liquids, chew gum or tobacco, or suck on hard candies for at least 3 hours prior to your arrival time. Do not smoke anything the morning of your procedure. ❑ Please take your medications as prescribed EXCEPT for those listed below. Medicines must be taken at least 3 hours prior to your arrival time. Take insulin according to PCP or endocrinologist instructions. ❑ Do not take diuretics (water/fluid pills), oral diabetic medications, iron pills, vitamins, or Carafate (sucralfate). If you have questions about what to take, please call us at least 1 business day before your arrival time. ❑ Bring a list of your current medications to your procedure.